

Polymorphisms in *pfert* and *pfmdr1*: parasite risk factors that affect treatment outcomes for falciparum malaria after artemether-lumefantrine and artesunate-amodiaquine

Meera Venkatesan^{1,2}, Nahla B. Gadalla³, Kasia Stepniewska⁴, Prabin Dahal⁴, Christian Nsanzabana⁴, Clarissa Moriera⁴, Ric N. Price^{4,5}, Andreas Mårtensson^{6,7}, Philip Rosenthal⁸, Grant Dorsey⁸, Colin J. Sutherland⁹, Philippe Guérin⁴, Timothy M. E. Davis¹⁰, Didier Ménard¹¹, Ishag Adam¹², George Ademowo¹³, Cesar Arze^{1,14}, Frederick N. Baliraine¹⁵, Nicole Berens-Riha¹⁶, Anders Björkman⁶, Steffen Borrmann^{17,18}, Francesco Checchi¹⁹, Meghna Desai²⁰, Mehul Dhorda^{1,21}, Abdoulaye A. Djimé²², Badria B. El-Sayed²³, Teferi Eshetu²⁴, Frederick Eyase²⁵, Catherine Falade²⁶, Jean-François Faucher^{27,28}, Gabrielle Fröberg⁶, Anastasia Grivoyannis²⁹, Sally Hamour³⁰, Sandrine Houzé³¹, Jacob Johnson³², Erasmus Kamugisha³³, Simon Kariuki, Jean-René Kiechel³⁴, Fred Kironde^{35,36}, Poul-Erik Kofoed⁶, Jacques LeBras^{31,37,38}, Maja Malmberg⁶, Leah Mwai³⁹, Billy Ngasala⁴⁰, Francois Nosten^{41,42}, Samuel L. Nsohya⁴³, Alexis Nzila⁴⁴, Mary Oguike⁹, Sabina Dahlström Otienoburu⁴⁵, Bernhards Ogutu³², Jean-Bosco Ouédraogo⁴⁶, Patrice Piola⁴⁷, Lars Rombo⁶, Birgit Schramm⁴⁸, A. Fabrice Somé⁴⁶, Julie Thwing⁵⁰, Johan Ursing⁶, Rina P. M. Wong¹⁰, Ahmed Zeynudin²⁴, Issaka Zongo⁴⁶, Carol Hopkins Sibley^{4,51*}, Christopher V. Plowe^{1,2}

- ¹ WorldWide Antimalarial Resistance Network Molecular Module (WWARN), University of Maryland School of Medicine, Baltimore, MD;
- ² Howard Hughes Medical Institute/Center for Vaccine Development, University of Maryland School of Medicine, Baltimore, MD;
- ³ National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD;
- ⁴ World Wide Antimalarial Resistance Network (WWARN), Centre for Tropical Medicine, Nuffield Department of Clinical Medicine, University of Oxford, Oxford, UK;
- ⁵ Global Health Division, Menzies School of Health Research and Charles Darwin University, Darwin, Australia;
- ⁶ Malaria Research, Infectious Diseases Unit, Department of Medicine Solna, Stockholm, Sweden;
- ⁷ Global Health, Department of Public Health Sciences, Karolinska Institutet, Stockholm, Sweden;
- ⁸ Department of Medicine, University of California, San Francisco, San Francisco, CA;
- ⁹ Department of Immunology and Infection, Faculty of Infectious and Tropical Diseases, London School of Hygiene and Tropical Medicine, London, UK;
- ¹⁰ School of Medicine and Pharmacology, The University of Western Australia, Fremantle Hospital, Nedlands, Australia;
- ¹¹ Malaria Molecular Epidemiology Unit, Institut Pasteur du Cambodge, Phnom Penh, Cambodia;
- ¹² Faculty of Medicine, University of Khartoum, Khartoum, Sudan;
- ¹³ Institute for Advanced Medical Research & Training, College of Medicine,, University of Ibadan;
- ¹⁴ Institute for Genome Sciences, University of Maryland School of Medicine, Baltimore, MD;
- ¹⁵ Department of Biology, LeTourneau University, Longview, TX;

- 16 Department of Infectious Diseases and Tropical Medicine, Ludwig Maximilians University, Munich, Germany;
- 17 Kenya Medical Research Institute/Wellcome Trust Research Programme, Kilifi, Kenya;
- 18 Institute of Tropical Medicine, University of Tübingen, Tübingen, Germany;
- 19 Save the Children, Paris, France;
- 20 Malaria Branch, Division of Parasitic Diseases, National Center for Zoonotic Vector Borne and Enteric Diseases, Centers for Disease Control and Prevention, Atlanta, GA;
- 21 Epicentre Uganda Research Base, Mbarara, Uganda;
- 22 Malaria Research and Training Centre, Faculty of Pharmacy, University of Science, Techniques and Technologies of Bamako, Bamako, Mali;
- 23 Department of Epidemiology, Tropical Medicine Research Institute, Khartoum, Sudan;
- 24 Department of Medical Laboratory Sciences and Pathology, Medical Parasitology Unit, Jimma University, Jimma, Ethiopia;
- 25 Global Emerging Infections System (GEIS), United States Army Research Unit-Kenya Walter Reed/KEMRI project, Kisumu, Kenya;
- 26 Department of Pharmacology & Therapeutics, University of Ibadan, Ibadan, Nigeria;
- 27 Department of Infectious Diseases, University Medical Center, Besançon, France;
- 28 Institut de Recherche pour le Développement (IRD), UMR 216, Paris, France ;
- 29 Department of Medicine, Division of Emergency Medicine, University of Washington, Seattle, WA;
- 30 UCL Centre for Nephrology, Royal Free Hospital, London, UK;
- 31 Laboratory of Parasitology, Malaria National Reference Centre, AP-HP Bichât Hospital, Paris France;
- 32 Institut de Médecine et d'Epidémiologie Appliquée, Bichât-C. Bernard Hospital, Paris, France;
- 33 United States Army Medical Research Unit-Kenya;
- 34 Catholic University of Health and Allied Sciences-Bugando, Mwanza, Tanzania;
- 35 Drugs for Neglected Diseases initiative (DNDi), Geneva, Switzerland;
- 36 Makerere University College of Health Sciences, Kampala, Uganda;
- 37 St. Augustine International University, Kampala, Uganda;
- 38 Institut de Recherche pour le Développement (IRD), UMR 216 Mother and Child face to Tropical Infections Research Unit, Paris, France;
- 39 PRES Paris Sorbonne Cité, Paris Descartes University, Paris, France;
- 40 Division of Translational Therapeutics, Department of Paediatrics, Faculty of Medicine, University of British Columbia, Vancouver, Canada;
- 41 Department of Parasitology, Muhimbili University of Health and Allied Sciences, Dar es Salaam, Tanzania;
- 42 Centre for Tropical Medicine, Nuffield Department of Medicine, University of Oxford, Oxford, UK;
- 43 Shoklo Malaria Research Unit, Mahidol-Oxford Tropical Medicine Research Unit, Faculty of Tropical Medicine, Mahidol University, Bangkok, Thailand;
- 44 Department of Pathology, School of Biomedical Science, Makerere University College of Health Sciences, Kampala, Uganda;
- 45 Department of Biology, King Fahd University of Petroleum and Minerals, Dharan, Saudi Arabia;
- 46 Institut de Médecine et d'Epidémiologie Appliquée, Bichât-C. Bernard Hospital;Worldwide Antimalarial Resistance Network (WWARN), Paris, France;

- ⁴⁷ Institut de Recherche en Sciences de la Santé, Centre Muraz, Bobo-Dioulasso, Burkina Faso;
⁴⁸ Institut Pasteur du Madagascar, Antananarivo, Madagascar;
⁴⁹ Epicentre, Paris, France
⁵⁰ Malaria Branch, Division of Parasitic Diseases and Malaria, Center for Global Health Centers for Disease Control and Prevention, Atlanta, GA;
⁵¹ Department of Genome Sciences, University of Washington, Seattle, WA

*corresponding author, Carol Hopkins Sibley, <carol.sibley@wwarn.org>

1 **Abstract**

2 Adequate clinical and parasitological cure by Artemisinin Combination Therapies (ACTs) relies
3 on both the artemisinin component and the partner drug. Polymorphisms in *pfcr* and *pfmdr1* are
4 associated with decreased sensitivity to amodiaquine and lumefantrine, but effects of these
5 polymorphisms on therapeutic responses to artesunate-amodiaquine and artemether-lumefantrine
6 have not been clearly defined. Individual patient data from 30 clinical trials were harmonized and
7 pooled, using standardized methodology from the WorldWide Antimalarial Resistance Network.
8 Data from more than 7000 patients were analyzed to assess relationships between parasite
9 polymorphisms in *pfcr* and *pfmdr1* and clinically relevant outcomes after treatment with AL or
10 ASAQ. Presence of *pfmdr1* N86 was a significant risk factors for recrudescence in patients treated
11 with AL, but elevated *pfmdr1* copy number was not. AL and ASAQ exerted opposing selective
12 effects on SNPs in *pfcr* and *pfmdr1*. Monitoring selection and responding to emerging signs of
13 drug resistance are critical tools for preserving ACT efficacy.
14
15

16 Introduction

17 Recent successes in malaria control have depended on the use of highly efficacious artemisinin
18 combination therapies (ACTs) for first-line treatment of uncomplicated *Plasmodium falciparum*
19 malaria. Adequate clinical and parasitological cure by ACTs relies on the rapid reduction in
20 parasite biomass by the potent, short-acting artemisinin component¹⁻³ and the subsequent
21 elimination of residual parasites by the longer-acting partner drug. The two most commonly used
22 ACTs worldwide are artemether-lumefantrine (AL) and artesunate-amodiaquine (ASAQ).⁴ PCR-
23 adjusted efficacy for both combinations remains high in most regions.⁵⁻⁷ However, there have been
24 some reports of decreasing AL cure rates in Africa⁸⁻¹¹ and Asia¹², and reports of high levels of
25 treatment failures of ASAQ.¹³⁻¹⁸ Resistance to ACT partner drugs has historically manifested
26 before that of artemisinins, whose short half-lives result in the exposure of residual parasites to
27 sub-therapeutic levels of the partner drug alone. Response to the partner drug is therefore a key
28 component of overall ACT efficacy.

29
30 Mutations in the gene encoding the *P. falciparum* chloroquine resistance transporter (*Pfcr*) are
31 associated with chloroquine resistance¹⁹; a change from lysine to threonine at codon 76 in *pfcr*
32 predicts responses of parasites to chloroquine.^{20, 21} In the presence of *pfcr* 76T, chloroquine
33 resistance is modulated by point mutations in the gene that encodes the *P. falciparum* multi drug
34 resistance transporter 1 (*pfmdr1*), primarily at codon 86^{22, 23} and also by mutations at positions
35 1034, 1042 and 1246.²⁴ Decreased susceptibility to lumefantrine has been linked to
36 polymorphisms in these two genes.²⁵⁻³⁵ Elevated *pfmdr1* copy number, which confers resistance
37 to mefloquine³⁶, has also been associated with reduced susceptibility to lumefantrine.³⁷⁻⁴⁰

38
39 Studies of amodiaquine have demonstrated reduced *in vivo* response⁴¹⁻⁴³ and increased IC₅₀ values
40 *in vitro*, in association with the presence of both *pfmdr1* 86Y and *pfcr* 76T alleles.^{44,45} Selection
41 of these alleles in recurrent parasites after treatment with amodiaquine alone or in combination
42 with artesunate has been observed in a number of studies.^{28,46-51} It has also been suggested that
43 parasites that carry chloroquine-resistant *pfmdr1* alleles may be more susceptible to artesunate in
44 classical *in vitro* assays,^{24,52} an effect that could counteract the increased risk of amodiaquine
45 failure when these drugs are combined in ASAQ.

46 Currently both AL and ASAQ retain high clinical efficacy with few recrudescence infections, and
47 individual studies generally lack sufficient statistical power to assess the association between
48 parasite genotypes and outcomes of clinical treatment. Such an assessment is a critical step in
49 validating molecular changes in parasite populations as useful markers of early signs of changing
50 parasite susceptibility to lumefantrine or amodiaquine. To overcome these challenges, individual
51 patient data on *in vivo* antimalarial efficacy and molecular markers of *P. falciparum* from 30
52 clinical trials were standardized, pooled, and nearly 7,000 patient responses were analyzed to
53 determine whether patients infected with parasites that carry these polymorphisms are at increased
54 risk of treatment failure. This large data set also provided the opportunity to examine the effects
55 of AL and ASAQ treatment on selection in parasites of particular alleles of *pfcr* and *pfmdr1*.

56 Methods

57 Selection and inclusion of data

58 Prospective clinical efficacy studies of *P. falciparum* treatment with AL (6 dose regimen) or
59 ASAQ (3 day fixed dose or loose/co-blistered regimen) with a minimum of 28 days of follow up
60 and genotyping of *pfcr* and/or *pfmdr1* were sought for the analysis. Studies were identified by a
61

62 systematic PubMed literature review using the search terms (artesunate AND amodiaquine) OR
63 (artemether AND lumefantrine) OR (ACT) AND (*pfmdr1* OR *pfprt*). Abstracts and text were
64 screened to determine whether inclusion criteria were met. Unpublished datasets were also
65 solicited and included in the analysis. Individual anonymized patient data including baseline
66 characteristics, drug intake, parasite density and temperature were collected. All but one study
67 included parasite genotyping to identify recrudescence infections of *P. falciparum*, and all studies
68 assessed the presence of *pfprt* and/or *pfmdr1* polymorphisms (single nucleotide polymorphisms
69 (SNPs) and copy number variation) in parasites isolated from patients on day 0. Multiplicity of
70 infection and molecular resistance marker data from other days including the day of microscopic
71 recurrent parasitemia were included but were not a prerequisite for study inclusion. Metadata on
72 study location, study design, drugs, and dosing regimens were also gathered. Figure 1 is a
73 schematic of the patient numbers and overall flow of the study.

74

75 ***Data curation and generation of variables***

76 All data sets were uploaded to the WWARN repository and standardized using the WWARN Data
77 Management and Statistical Analysis Plans (DMSAP).^{53,54} Outcome status and censoring were
78 defined according to the Clinical DMSAP.⁵³ Parasites that recurred within the follow-up period
79 were classified using WHO guidelines⁵⁵: microscopically-detected infections during follow-up
80 were classified as ‘recurrent’; recurrent infections sharing with blood samples taken at day 0 PCR
81 bands in polymorphic merozoite antigens or microsatellite fragment sizes were termed
82 ‘recrudescence’, and recurrent infections not sharing PCR bands or microsatellite fragment sizes
83 with blood samples taken at day 0 were classified as ‘re-infections’ (new infections). Molecular
84 markers were coded as either single or mixed allele genotypes in the case of SNPs and as mean
85 copy number per sample for copy number polymorphisms. Multi-SNP haplotypes were
86 reconstructed as described in the Molecular DMSAP.^{56,57}

87 ***Statistical analysis***

88 All statistical analyses were conducted using Stata 11 (Stata Corp, College Station, Texas). The
89 primary endpoint was clinical efficacy, defined as the PCR-adjusted risk of *P. falciparum*
90 recrudescence infections. The cumulative risk of recrudescence at day 28 and day 42 was computed
91 using survival analysis [Kaplan-Meier estimates (K-M)]. Comparisons of K-M survival curves
92 were performed using log rank tests stratified by study sites.

93

94 Multivariable analysis of risk factors associated with PCR-adjusted recrudescence was conducted
95 using Cox proportional hazards regression models with shared frailty parameters to adjust for site-
96 specific effects. The risk factors that affect the clinical efficacy of AL and ASAQ have been
97 intensively studied in pooled analyses of both ACTs. Sixty-two studies with 14,651 patients
98 treated with AL and 39 studies including 8,337 patients treated with ASAQ were analyzed; these
99 full analyses have been submitted for publication. The univariable and multivariable risk factors
100 identified in those studies are shown in supplementary tables S1A and S1B. Clinical covariates in
101 the current study were included based on the previous analyses as follows: (lumefantrine or
102 amodiaquine dose, enrolment parasitemia, age category, and ASAQ fixed or co-blistered versus
103 loose formulation (Table 1). Each molecular marker was then added to the model. The proportional
104 hazard assumption was tested based on Schoenfeld residuals⁵⁸. In the case of non-proportionality,
105 interactions with a categorized time variable based on clinical follow-up intervals (< day 14, days
106 14 – 21, 21 – 28, and > day 28) were used to account for changing effects over time, and
107 neighboring windows with similar effects of genetic covariates as determined by Wald test were

108 merged. Finally, other covariates (transmission intensity, region of sample origin, dose
109 supervision, and fat intake) were included in the model if they improved model fit based on the
110 likelihood ratio test. Multiplicity of infection was only available for 197 and 141 AL and ASAQ
111 patients, respectively, and was excluded from further analysis. The final model was then used to
112 estimate the adjusted hazard ratio for recrudescence in patients who carried parasites with resistant
113 versus sensitive genotypes on day 0. The assumption of proportional hazards was tested separately
114 for the individual covariates in the final multivariable model, and any violations were reported.
115

116 In patients who had recurrent parasitemia on or before day 42, changes in *pfprt* and *pfmdr1* alleles
117 between pre- and post-treatment matched pairs of samples was compared using McNemar's test.
118 Changes in genotype, rather than presence of a particular allele, were compared between matched
119 pairs to ensure that differences reflected selection rather than underlying differences in allele
120 frequencies among populations. The effect of markers present at the time of recurrence on median
121 time to PCR-adjusted re-infection (new infection) was investigated using the Wilcoxon Mann-
122 Whitney U test. Competing risk analysis⁵⁹ was used to estimate cumulative incidence of PCR-
123 adjusted re-infections with specific genotypes, where recrudescence and re-infections with other
124 genotypes were treated as competing events.
125

126 The number of molecular markers used to distinguish recrudescence from re-infection varied from
127 one to three or more loci. The effect of the number of loci genotyped on outcome classification
128 was investigated in a regression model of predictors of recrudescence within all recurrences. No
129 effect of this variable was observed on the number of recrudescence infections identified among
130 recurrences in univariable or multivariable analysis, it was not further investigated.
131

132 **Results**

133 Individual patient and linked parasite genotype data from 30 studies were available (Listed
134 individually in Supplementary Table 2). Data from 6,947 patients who were treated with AL
135 (4,701) or ASAQ (2,246) were included in the analysis. Twenty two studies were published,
136 representing 91% of all published clinical data on AL and ASAQ in which *pfprt* or *pfmdr1*
137 genotypes were determined. Baseline characteristics for patients treated with AL or ASAQ are
138 presented in Supplementary Table S3.
139

140 ***Clinical efficacy of AL and ASAQ***

141 The estimates of efficacy (defined as risk of PCR-adjusted recrudescence) of AL and ASAQ are
142 summarized in Table 2. Of the 4,701 AL patients, 4,504 were followed up for at least one day and
143 were included in the analysis. Similarly, of the 2,246 ASAQ patients, 2,099 were included. In total,
144 1,051 patients had recurrent parasitemia following AL, of which 155 (15%) were classified by
145 PCR as recrudescence infections. The corresponding figures for ASAQ were 484 patients had
146 recurrent parasitemia and 58 (12%) were confirmed as recrudescence. The overall clinical efficacy
147 at day 42 was 95.3% [95%CI: 94.4-96.0] in patients treated with AL and 95.1% [95%CI: 92.3-
148 96.7] following ASAQ treatment (Table 2). The proportion of adequate clinical and parasitological
149 response of ASAQ was significantly higher for the fixed dose and co-blistered tablets (97.0% [95%
150 CI: 94.4-98.4] compared to the loose formulation (93.0% [95% CI: 89.2-95.6] (p=0.003).
151

152 ***Baseline prevalence of genetic markers associated with resistance***

153 The baseline prevalence of SNPs in *pfprt* and *pfmdr1* was determined, but not all SNPs were
154 available for all isolates. The most frequently analyzed SNPs were position 76 in *pfprt* determined
155 in 3,640 patients and position 86 in *pfmdr1* in 3,510, with the complete haplotype of positions 72-
156 76 in *pfprt*, *pfmdr1* copy number, and SNPs at positions *pfmdr1* 184 and and 1246 available in a
157 subset of patients (Table 3).

158
159 The prevalence of *pfprt* and *pfmdr1* alleles varied by region (Table 3). The *pfprt* 76T allele (all in
160 the SVMNT haplotype) was almost fixed at 96.4% (81/84) in isolates from Asia (Thailand) and
161 Oceania (Papua New Guinea). In Africa, the only resistant haplotype observed was the CVIET
162 allele. The 76T allele predominated: 67.6% (1155/1708) in East Africa and 73.3% (1,354/1,848)
163 in West Africa (Table 3). Amplification of *pfmdr1* was seen in 51% (54/106) of Asian isolates
164 examined for this genotype, but only in 2.4% (17/659) of isolates from Africa. *Pfmdr1* 86Y was
165 found in 38% (59/156) of isolates from Asia/Oceania; in contrast, the 86Y allele was present in
166 44% (896/2,033) of isolates from East Africa and 34.3% (453/1,321) of isolates from West Africa.

167
168 The SNPs at positions 184 and 1246 showed similar patterns, with *pfmdr1* Y184 and D1246
169 predominating in all three regions (Table 3). Almost all isolates examined carried the *pfmdr1*
170 S1034 (703/786) and N1042 (997/1,005).

171
172 ***Parasite genotypes as risk factors for recrudescence infection***
173 After controlling for age, baseline parasite density, and total lumefantrine dose (Table 1), the
174 presence of parasites in the initial infection that carried *pfmdr1* N86 (alone or a mixed infection
175 with *pfmdr1* 86Y) was a significant risk factor for recrudescence infection occurring between days
176 14 and 28 after AL treatment (Adjusted Hazards Ratio AHR = 4.06 (95% CI [1.94– 8.47]; $p <$
177 0.001) (Table 4, Figure 2A). Region of sample origin was not included as a covariate in the model
178 because it violated the assumption of proportional hazards.

179
180 The presence of more than one copy of *pfmdr1* was not a significant risk factor for recrudescence
181 after AL treatment and did not become a significant risk factor when the effect of region was added
182 to the model. Once adjusted by study site, AL efficacy in patients with infections with a single
183 copy of *pfmdr1* was not significantly different from efficacy in patients with multi-copy infections
184 ($p=0.47$; Figure 2B). The interaction of region of origin with *pfmdr1* copy number could not be
185 investigated because of insufficient multi-copy samples from Africa in the model. Because *pfmdr1*
186 copy number was not found to be a significant risk factor for AL recrudescence, the model
187 including both *pfmdr1* 86 and *pfmdr1* copy number is not included here or presented in Table 4.

188 No association was observed between the *pfmdr1* 184, *pfmdr1* 1246, and *pfprt* polymorphisms and
189 recrudescence infections after AL treatment. The risk for parasites with the *pfmdr1* N86 + D1246
190 haplotype is not reported here because it represents a subset of the *pfmdr1* N86 sample set (of the
191 samples genotyped for both SNPs, all but 17 samples with *pfmdr1* N86 also had D1246).

192
193 For patients treated with ASAQ, none of the analyzed *pfprt* or *pfmdr1* parasite genotypes were
194 significant risk factors for recrudescence infections in the multivariable analysis.

195
196 ***Post-treatment selection of genetic markers associated with resistance***

197 To examine changes in the genotypes of parasites following drug treatment, we compared the
198 prevalence of *pfmdr1* and *pfcr1* alleles in paired isolates from the initial and the recurrent parasites
199 in the subset of patients in whom parasites recurred during the 42 day follow up period. Post-
200 treatment changes among specific genotypes are presented in Table 5 for all recurrent infections.
201 Significant selection of *pfcr1* K76 and *pfmdr1* N86 occurred in both recrudescence and re-infecting
202 parasites after AL treatment. Selection of *pfmdr1* 184F and D1246 alleles was also observed in
203 the recurrent parasites and *pfmdr1* D1246 in those that reinfected patients after treatment. Selection
204 of single or multiple copies of *pfmdr1* was not observed in any of the groups (Table 5). *Pfmdr1*
205 86Y and 1246Y were significantly selected in recurrent and re-infections after treatment with
206 ASAQ (Table 5).

207

208 **Median time to re-infection**

209 The genotype of parasites at the time of re-infection provides another metric of their susceptibility
210 to a drug. This analysis indicated that in patients treated with AL, re-infecting parasites carrying
211 *pfmdr1* N86, *pfmdr1* D1246 or *pfcr1* K76 alleles appeared earlier than those carrying *pfmdr1* 86Y,
212 *pfmdr1* 1246Y or *pfcr1* 76T (Figure 3A). Correspondingly, in patients treated with AL, parasites
213 carrying *pfmdr1* N86 had a median time to re-infection of 28 days (interquartile range = 21–35)
214 compared to 35 days (interquartile range = 28-42) for those with *pfmdr1* 86Y ($p < 0.001$). Similar
215 differences in the time to re-infection were observed for patients infected with parasites that carried
216 the *pfmdr1* 184F ($p = 0.008$) or *pfcr1* K76 alleles ($p = 0.001$) when compared to *pfmdr1* Y184 or *pfcr1*
217 76T.

218

219 In contrast, in patients treated with ASAQ, parasites carrying *pfmdr1* 86Y, *pfmdr1* 1246Y, or *pfcr1*
220 76T appeared earlier after treatment than those carrying *pfmdr1* N86, *pfmdr1* D1246 or *pfcr1* K76
221 (Figure 3B). Parasites with *pfcr1* 76T had a median re-infection day of 28 (interquartile range =
222 21–35) compared to day 37.5 (interquartile range = 28-42) for those carrying K76 ($p = 0.053$) and
223 those with *pfmdr1* 1246Y re-infected on a median day of 21 (interquartile range = 21-28) compared
224 to day 28 (interquartile range = 21-35) for those with D1246 ($p = 0.001$).

225

226 **Discussion**

227 This pooled analysis of data from 31 clinical studies shows clearly that the genotypes of infecting
228 parasites influence the outcome of AL treatment. Patients infected with parasites that carried the
229 *pfmdr1* N86 allele or increased *pfmdr1* copy number were at significantly greater risk of treatment
230 failure than those whose parasites carried the 86Y allele or a single copy of *pfmdr1*. Analysis of
231 the clinical outcomes after treatment with ASAQ did not link a particular genotype with treatment
232 failure in this smaller data set, however it did show clear evidence of selection of particular parasite
233 genotypes. Our findings are consistent with previous molecular studies in which changes in the
234 prevalence of particular alleles of *pfcr1* or *pfmdr1* have been documented in response to
235 introduction or increased use of lumefantrine²⁵⁻³⁵ or amodiaquine.^{15, 28, 40-51}

236

237 Our observation that parasites with the *pfmdr1* N86, D1246, and *pfcr1* K76 alleles re-infected
238 patients earlier after AL treatment, and parasites carrying the *pfmdr1* 86Y, 1246Y, and *pfcr1* 76T
239 alleles re-infected patients earlier after ASAQ is also congruent with the molecular studies. These
240 differences suggest that parasites with these genotypes can withstand higher drug concentrations
241 compared with parasites that carry the alternative alleles. Recently, Malmberg and colleagues
242 demonstrated this effect quantitatively. After AL treatment, parasites with the *pfmdr1*

243 N86/184F/D1246 haplotype were able to re-infect patients whose lumefantrine blood
244 concentrations were 15-fold higher than was the case for parasites carrying the 86Y/Y184/1246Y
245 haplotype,³³ providing a potential pharmacological explanation for the molecular findings.
246 Together, these observations suggest that monitoring shifts to earlier time of re-infection could
247 provide a relatively simple warning of declining susceptibility to these drugs, especially if
248 combined with timed measurement of drug concentrations in patients' blood.

249 In Southeast Asia, parasites with multi-copy *pfmdr1* are common in areas where mefloquine has
250 been intensively deployed,³⁶ and increased *pfmdr1* copy number is strongly associated with
251 artesunate-mefloquine treatment failures. Almost half of the samples in our data set from Southeast
252 Asia region had at least two copies of the gene. In contrast, multiple copy number was rarely
253 observed in our large sample of isolates from Africa, where populations have had little exposure
254 to mefloquine.

255 The results of this study did not indicate that parasites with increased copy number of *pfmdr1* are
256 less sensitive to lumefantrine. Our findings contrast with reports of decreased *in vitro* lumefantrine
257 susceptibility with increased copy number³⁷⁻⁴⁰, but support the conclusions of *in vivo* studies³⁸
258 which indicate that multi-copy *pfmdr1* is not a risk factor for AL treatment failure.

259 In our data set from Southeast Asia, the amplified alleles all carried the N86 allele of *pfmdr1*^{34,36,62}.
260 This association was not found in the few parasites from Africa in our data set that did have an
261 increased copy number³¹, indicating that either of the N86Y alleles of *pfmdr1* can apparently be
262 amplified in this region.

263 The evidence of strong selection of particular alleles by both drugs in recurrent parasites, coupled
264 with our observation that particular parasite genotypes increase risk of treatment failure,
265 demonstrates that tracking these molecular markers can signal early declines in susceptibility to
266 lumefantrine or amodiaquine. Both alleles of *pfmdr1* N86Y, Y184F and D1246Y are common in
267 African *P. falciparum* populations, and *pfprt* K76 has increased in prevalence in recent years, so
268 changes in the prevalence of these alleles can be a sensitive indicator of selection of parasite
269 populations by AL and ASAQ. In turn, declining efficacy of these partner drugs exposes the
270 artemether or artesunate component of the ACT to selective pressure and could facilitate
271 emergence of new foci of resistance to artemisinin, as observed in the Mekong region. The recent
272 identification of a marker correlated with slow response to artemisinin,⁶³ will allow molecular
273 assessment of this trend, as well.

274
275 Application of these molecular tools is increasingly feasible both in the context of clinical trials
276 and in community surveys of populations where AL or ASAQ are heavily used. These approaches
277 can offer cost effective methods that detect evidence of declines in parasite susceptibility far earlier
278 than before, allowing detailed studies of clinical responses to the drugs in areas of concern. This
279 two stage approach can provide an opportunity for policy makers to manage emerging threats of
280 resistance before clinical failure of a drug is manifest and preserve the useful therapeutic life of
281 these valuable antimalarials for as long as possible.

282
283 Finally, these results suggest that both AL and ASAQ interact with the proteins encoded by *pfprt*
284 and *pfmdr1*, but the two drugs select alternative alleles. This opposing selection of parasite
285 genotypes by the partner drugs could influence the choice of an ACT in regions with different

286 patterns of *pfprt* and *pfmdr1* polymorphisms. For example, if a particular allele is rapidly
287 increasing under intensive use of AL, introduction of AQ might be introduced to counteract that
288 trend. Concurrent use of these two ACTs that exert opposing selective pressures on recurrent
289 parasites could provide a counterbalance and prevent strong directional selection in both *pfprt* and
290 *pfmdr1*, maintaining the overall efficacy of both AL and ASAQ for a long period. Despite logistical
291 challenges, the simultaneous use of multiple first line therapies is supported by mathematical
292 models,⁶⁴⁻⁶⁶ and concurrent availability of AL and ASAQ, as implemented in some West African
293 countries⁴ may provide a practical means to test this strategy directly.

294

295 *Acknowledgements*

296 The WorldWide Antimalarial Resistance Network is supported by the Bill & Melinda Gates
297 Foundation. The opinions and assertions contained herein are the personal opinions of the authors
298 (JJ and FE) and are not to be construed as reflecting the views of the US Army Medical Research
299 Unit-Kenya or the US Department of Defense. The authors would like to thank Dr. Pascal
300 Ringwald for his review of the *pfmdr1* copy number results which led to the corrections published
301 in 2018.

302

303

References

1. Bethell DB, Teja-Isavadharm P, Cao XT, Pham TT, Ta TT, Tran TN, Nguyen TT, Pham TP, Kyle D, Day NP, White NJ, 1997. Pharmacokinetics of oral artesunate in children with moderately severe *Plasmodium falciparum* malaria. *Trans.R.Soc.Trop.Med.Hyg.* 91: 195-198.
2. Na BK, Karbwang J, Thomas CG, Thanavibul A, Sukontason K, Ward SA, Edwards G, 1994. Pharmacokinetics of artemether after oral administration to healthy Thai males and patients with acute, uncomplicated falciparum malaria. *Br.J.Clin.Pharmacol.* 37: 249-253.
3. White NJ, van VM, Ezzet F, 1999. Clinical pharmacokinetics and pharmacodynamics of artemether-lumefantrine. *Clin.Pharmacokinet.* 37: 105-125.
4. Organization WH, 2013. World Malaria Report 2013. Available at: http://www.who.int/malaria/publications/world_malaria_report_2013/report/en/. Accessed December 30, 2013, 2013.
5. Makanga M, Krudsood S, 2009. The clinical efficacy of artemether/lumefantrine (Coartem). *Malar.J.* 8 Suppl 1: S5.
6. Sinclair D, Zani B, Donegan S, Olliaro P, Garner P, 2009. Artemisinin-based combination therapy for treating uncomplicated malaria. *Cochrane.Database.Syst.Rev.:* CD007483.
7. Stover KR, King ST, Robinson J, 2012. Artemether-lumefantrine: an option for malaria. *Ann.Pharmacother.* 46: 567-577.
8. Mukhtar EA, Gadalla NB, El-Zaki SE, Mukhtar I, Mansour FA, Babiker A, El-Sayed BB, 2007. A comparative study on the efficacy of artesunate plus sulphadoxine/pyrimethamine versus artemether-lumefantrine in eastern Sudan. *Malar.J.* 6: 92.
9. Siribie M, Diarra A, Tiono AB, Soulama I, Sirima SB, 2012. [Efficacy of artemether-lumefantrine in the treatment of uncomplicated malaria in children living in a rural area of Burkina Faso in 2009]. *Bull.Soc.Pathol.Exot.* 105: 202-207.
10. Abuaku B, Duah N, Quaye L, Quashie N, Koram K, 2012. Therapeutic efficacy of artemether-lumefantrine combination in the treatment of uncomplicated malaria among children under five years of age in three ecological zones in Ghana. *Malar.J.* 11: 388.
11. Ngasala BE, Malmberg M, Carlsson AM, Ferreira PE, Petzold MG, Blessborn D, Bergqvist Y, Gil JP, Premji Z, Mårtensson A, 2011. Effectiveness of artemether-lumefantrine provided by community health workers in under-five children with uncomplicated malaria in rural Tanzania: an open label prospective study. *Malar.J.* 10: 64.
12. Song J, Socheat D, Tan B, Seila S, Xu Y, Ou F, Sokunthea S, Sophorn L, Zhou C, Deng C, Wang Q, Li G, 2011. Randomized trials of artemisinin-piperaquine, dihydroartemisinin-piperaquine phosphate and artemether-lumefantrine for the treatment of multi-drug resistant falciparum malaria in Cambodia-Thailand border area. *Malar.J.* 10: 231.
13. Mutabingwa TK, Anthony D, Heller A, Hallett R, Ahmed J, Drakeley C, Greenwood BM, Whitty CJ, 2005. Amodiaquine alone, amodiaquine+sulfadoxine-pyrimethamine, amodiaquine+artesunate, and artemether-lumefantrine for outpatient treatment of malaria in Tanzanian children: a four-arm randomised effectiveness trial. *Lancet* 365: 1474-1480.
14. Mårtensson A, Stromberg J, Sisowath C, Msellem MI, Gil JP, Montgomery SM, Olliaro P, Ali AS, Bjorkman A, 2005. Efficacy of artesunate plus amodiaquine versus that of artemether-lumefantrine for the treatment of uncomplicated childhood *Plasmodium falciparum* malaria in Zanzibar, Tanzania. *Clin.Infect.Dis.* 41: 1079-1086.
15. Thwing JI, Odero CO, Odhiambo FO, Otieno KO, Kariuki S, Ord R, Roper C, McMorro M, Vulule J, Slutsker L, Newman RD, Hamel MJ, Desai M, 2009. In-vivo efficacy of

- amodiaquine-artesunate in children with uncomplicated *Plasmodium falciparum* malaria in western Kenya. *Trop.Med.Int.Health* 14: 294-300.
16. Hamour S, Melaku Y, Keus K, Wambugu J, Atkin S, Montgomery J, Ford N, Hook C, Checchi F, 2005. Malaria in the Nuba Mountains of Sudan: baseline genotypic resistance and efficacy of the artesunate plus sulfadoxine-pyrimethamine and artesunate plus amodiaquine combinations. *Trans.R.Soc.Trop.Med.Hyg.* 99: 548-554.
 17. Swarthout TD, van den Broek IV, Kayembe G, Montgomery J, Pota H, Roper C, 2006. Artesunate + amodiaquine and artesunate + sulphadoxine-pyrimethamine for treatment of uncomplicated malaria in Democratic Republic of Congo: a clinical trial with determination of sulphadoxine and pyrimethamine-resistant haplotypes. *Trop.Med.Int.Health* 11: 1503-1511.
 18. Rwagacondo CE, Karema C, Mugisha V, Erhart A, Dujardin JC, Van OC, Ringwald P, D'Alessandro U, 2004. Is amodiaquine failing in Rwanda? Efficacy of amodiaquine alone and combined with artesunate in children with uncomplicated malaria. *Trop.Med.Int.Health* 9: 1091-1098.
 19. Fidock DA, Nomura T, Talley AK, Cooper RA, Dzekunov SM, Ferdig MT, Ursos LM, Sidhu AB, Naude B, Deitsch KW, Su XZ, Wootton JC, Roepe PD, Wellems TE, 2000. Mutations in the *P. falciparum* digestive vacuole transmembrane protein *Pfcr* and evidence for their role in chloroquine resistance. *Mol.Cell* 6: 861-871.
 20. Djimdé A, Doumbo OK, Cortese JF, Kayentao K, Doumbo S, Diourte Y, Coulibaly D, Dicko A, Su XZ, Nomura T, Fidock DA, Wellems TE, Plowe CV, 2001. A molecular marker for chloroquine-resistant falciparum malaria. *N.Engl.J.Med.* 344: 257-263.
 21. Djimdé AA, Barger B, Kone A, Beavogui AH, Tekete M, Fofana B, Dara A, Maiga H, Dembele D, Toure S, Dama S, Ouologuem D, Sangare CP, Dolo A, Sogoba N, Nimaga K, Kone Y, Doumbo OK, 2010. A molecular map of chloroquine resistance in Mali. *FEMS Immunol Med Microbiol* 58: 113-8.
 22. Babiker HA, Pringle SJ, Abdel-Muhsin A, Mackinnon M, Hunt P, Walliker D, 2001. High-level chloroquine resistance in Sudanese isolates of *Plasmodium falciparum* is associated with mutations in the chloroquine resistance transporter gene *pfcr* and the multidrug resistance Gene *pfmdr1*. *J.Infect.Dis.* 183: 1535-1538.
 23. Mu J, Ferdig MT, Feng X, Joy DA, Duan J, Furuya T, Subramanian G, Aravind L, Cooper RA, Wootton JC, Xiong M, Su XZ, 2003. Multiple transporters associated with malaria parasite responses to chloroquine and quinine. *Mol.Microbiol.* 49: 977-989.
 24. Reed MB, Saliba KJ, Caruana SR, Kirk K, Cowman AF, 2000. Pgh1 modulates sensitivity and resistance to multiple antimalarials in *Plasmodium falciparum*. *Nature* 403: 906-909.
 25. Sisowath C, Stromberg J, Mårtensson A, Msellem M, Obondo C, Bjorkman A, Gil JP, 2005. In vivo selection of *Plasmodium falciparum pfmdr1* 86N coding alleles by artemether-lumefantrine (Coartem). *J.Infect.Dis.* 191: 1014-1017.
 26. Dokomajilar C, Nsobya SL, Greenhouse B, Rosenthal PJ, Dorsey G, 2006. Selection of *Plasmodium falciparum pfmdr1* alleles following therapy with artemether-lumefantrine in an area of Uganda where malaria is highly endemic. *Antimicrob.Agents Chemother.* 50: 1893-1895.
 27. Sisowath C, Ferreira PE, Bustamante LY, Dahlstrom S, Mårtensson A, Bjorkman A, Krishna S, Gil JP, 2007. The role of *pfmdr1* in *Plasmodium falciparum* tolerance to artemether-lumefantrine in Africa. *Trop.Med.Int.Health* 12: 736-742.

28. Humphreys GS, Merinopoulos I, Ahmed J, Whitty CJ, Mutabingwa TK, Sutherland CJ, Hallett RL, 2007. Amodiaquine and artemether-lumefantrine select distinct alleles of the *Plasmodium falciparum* *mdr1* gene in Tanzanian children treated for uncomplicated malaria. *Antimicrob.Agents Chemother.* 51: 991-997.
29. Uhlemann AC, McGready R, Ashley EA, Brockman A, Singhasivanon P, Krishna S, White NJ, Nosten F, Price RN, 2007. Intra-host selection of *Plasmodium falciparum* *pfmdr1* alleles after antimalarial treatment on the northwestern border of Thailand. *J.Infect.Dis.* 195: 134-141.
30. Somé AF, Sere YY, Dokomajilar C, Zongo I, Rouamba N, Greenhouse B, Ouedraogo JB, Rosenthal PJ, 2010. Selection of known *Plasmodium falciparum* resistance-mediating polymorphisms by artemether-lumefantrine and amodiaquine-sulfadoxine-pyrimethamine but not dihydroartemisinin-piperaquine in Burkina Faso. *Antimicrob.Agents Chemother.* 54: 1949-1954.
31. Gadalla NB, Adam I, Elzaki SE, Bashir S, Mukhtar I, Oguike M, Gadalla A, Mansour F, Warhurst D, El-Sayed BB, Sutherland CJ, 2011. Increased *pfmdr1* copy number and sequence polymorphisms in *Plasmodium falciparum* isolates from Sudanese malaria patients treated with artemether-lumefantrine. *Antimicrob.Agents Chemother.* 55: 5408-5411.
32. Griyovannis A, Niangaly, Traore OB, Kodio A, Traore K, Tolo Y, Dembele A, Traore A, Bamadio A, Traore ZI, Sanogo K, Sissoko M, Sagara I, Djimdé AA, Doumbo OK, In vivo Selection of *Plasmodium falciparum* *pfert* K76 and *pfmdr1* N86 alleles by Artemether-Lumefantrine in Mali.
33. Malmberg M, Ferreira PE, Tarning J, Ursing J, Ngasala B, Bjorkman A, Mårtensson A, Gil JP, 2013. *Plasmodium falciparum* drug resistance phenotype as assessed by patient antimalarial drug levels and its association with *pfmdr1* polymorphisms. *J.Infect.Dis.* 207: 842-847.
34. Price RN, Uhlemann AC, van VM, Brockman A, Hutagalung R, Nair S, Nash D, Singhasivanon P, Anderson TJ, Krishna S, White NJ, Nosten F, 2006. Molecular and pharmacological determinants of the therapeutic response to artemether-lumefantrine in multidrug-resistant *Plasmodium falciparum* malaria. *Clin.Infect.Dis.* 42: 1570-1577.
35. Happi CT, Gbotosho GO, Folarin OA, Sowunmi A, Hudson T, O'Neil M, Milhous W, Wirth DF, Oduola AM, 2009. Selection of *Plasmodium falciparum* multidrug resistance gene 1 alleles in asexual stages and gametocytes by artemether-lumefantrine in Nigerian children with uncomplicated falciparum malaria. *Antimicrob.Agents Chemother.* 53: 888-895.
36. Price RN, Uhlemann AC, Brockman A, McGready R, Ashley E, Phaipun L, Patel R, Laing K, Looareesuwan S, White NJ, Nosten F, Krishna S, 2004. Mefloquine resistance in *Plasmodium falciparum* and increased *pfmdr1* gene copy number. *Lancet* 364: 438-447.
37. Sidhu AB, Uhlemann AC, Valderramos SG, Valderramos JC, Krishna S, Fidock DA, 2006. Decreasing *pfmdr1* copy number in *Plasmodium falciparum* malaria heightens susceptibility to mefloquine, lumefantrine, halofantrine, quinine, and artemisinin. *J.Infect.Dis.* 194: 528-535.
38. Lim P, Alker AP, Khim N, Shah NK, Incardona S, Doung S, Yi P, Bouth DM, Bouchier C, Puijalon OM, Meshnick SR, Wongsrichanalai C, Fandeur T, Le BJ, Ringwald P, Ariey F, 2009. *Pfmdr1* copy number and artemisinin derivatives combination therapy failure in falciparum malaria in Cambodia. *Malar.J.* 8: 11.

39. Mungthin M, Khositnithikul R, Sitthichot N, Suwandittakul N, Wattanaveeradej V, Ward SA, Na-Bangchang K, 2010. Association between the *pfmdr1* gene and in vitro artemether and lumefantrine sensitivity in Thai isolates of *Plasmodium falciparum*. *Am.J.Trop.Med.Hyg.* 83: 1005-1009.
40. Simpson JA, Jansen KM, Anderson TJ, Zaloumis S, Nair S, Woodrow C, White NJ, Nosten F, Price RN, 2013. Nonlinear Mixed-Effects Modelling of In Vitro Drug Susceptibility and Molecular Correlates of Multidrug Resistant *Plasmodium falciparum*. *PLoS.One.* 8: e69505.
41. Ochong EO, van den Broek IV, Keus K, Nzila A, 2003. Short report: association between chloroquine and amodiaquine resistance and allelic variation in the *Plasmodium falciparum* multiple drug resistance 1 gene and the chloroquine resistance transporter gene in isolates from the upper Nile in southern Sudan. *Am.J.Trop.Med.Hyg.* 69: 184-187.
42. Happi CT, Gbotosho GO, Folarin OA, Bolaji OM, Sowunmi A, Kyle DE, Milhous W, Wirth DF, Oduola AM, 2006. Association between mutations in *Plasmodium falciparum* chloroquine resistance transporter and *P. falciparum* multidrug resistance 1 genes and in vivo amodiaquine resistance in *P. falciparum* malaria-infected children in Nigeria. *Am.J.Trop.Med.Hyg.* 75: 155-161.
43. Tinto H, Guekoun L, Zongo I, Guiguemde RT, D'Alessandro U, Ouedraogo JB, 2008. Chloroquine-resistance molecular markers (*Pfcr1* T76 and *Pfmdr-1* Y86) and amodiaquine resistance in Burkina Faso. *Trop.Med.Int.Health* 13: 238-240.
44. Echeverry DF, Holmgren G, Murillo C, Higueta JC, Bjorkman A, Gil JP, Osorio L, 2007. Short report: polymorphisms in the *pfcr1* and *pfmdr1* genes of *Plasmodium falciparum* and in vitro susceptibility to amodiaquine and desethylamodiaquine. *Am.J.Trop.Med.Hyg.* 77: 1034-1038.
45. Folarin OA, Bustamante C, Gbotosho GO, Sowunmi A, Zalis MG, Oduola AM, Happi CT, 2011. In vitro amodiaquine resistance and its association with mutations in *pfcr1* and *pfmdr1* genes of *Plasmodium falciparum* isolates from Nigeria. *Acta Trop.* 120: 224-230.
46. Nsobya SL, Dokomajilar C, Joloba M, Dorsey G, Rosenthal PJ, 2007. Resistance-mediating *Plasmodium falciparum* *pfcr1* and *pfmdr1* alleles after treatment with artesunate-amodiaquine in Uganda. *Antimicrob.Agents Chemother.* 51: 3023-3025.
47. Holmgren G, Hamrin J, Svard J, Mårtensson A, Gil JP, Bjorkman A, 2007. Selection of *pfmdr1* mutations after amodiaquine monotherapy and amodiaquine plus artemisinin combination therapy in East Africa. *Infect.Genet.Evol.* 7: 562-569.
48. Holmgren G, Gil JP, Ferreira PM, Veiga MI, Obonyo CO, Bjorkman A, 2006. Amodiaquine resistant *Plasmodium falciparum* malaria in vivo is associated with selection of *pfcr1* 76T and *pfmdr1* 86Y. *Infect.Genet.Evol.* 6: 309-314.
49. Djimdé AA, Fofana B, Sagara I, Sidibe B, Toure S, Dembele D, Dama S, Ouologuem D, Dicko A, Doumbo OK, 2008. Efficacy, safety, and selection of molecular markers of drug resistance by two ACTs in Mali. *Am.J.Trop.Med.Hyg.* 78: 455-461.
50. Danquah I, Coulibaly B, Meissner P, Petruschke I, Muller O, Mockenhaupt FP, 2010. Selection of *pfmdr1* and *pfcr1* alleles in amodiaquine treatment failure in north-western Burkina Faso. *Acta Trop.* 114: 63-66.
51. Duraisingh MT, Drakeley CJ, Muller O, Bailey R, Snounou G, Targett GA, Greenwood BM, Warhurst DC, 1997. Evidence for selection for the tyrosine-86 allele of the *pfmdr1* gene of *Plasmodium falciparum* by chloroquine and amodiaquine. *Parasitology* 114 (Pt 3): 205-211.

52. Duraisingh MT, Jones P, Sambou I, von SL, Pinder M, Warhurst DC, 2000. The tyrosine-86 allele of the *pfmdr1* gene of *Plasmodium falciparum* is associated with increased sensitivity to the anti-malarials mefloquine and artemisinin. *Mol.Biochem.Parasitol.* 108: 13-23.
53. WWARN, 2012. Clinical Module: Data Management and Statistical Analysis Plan. Version 1.2. Available at: <http://www.wwarn.org/sites/default/files/ClinicalDMSAP.pdf>. Accessed Jan 13, 2014.
54. WWARN, 2012. Statistical Analysis Plan AL Dose Impact Study Group Version 1.9. Available at: <http://www.wwarn.org/sites/default/files/ALDoseImpactStudyGroupSAP.pdf>. Accessed Jan 13, 2014.
55. WHO, 2008. Methods and techniques for clinical trials on antimalarial drug efficacy: genotyping to identify parasite populations. Geneva: World Health Organization.
56. WWARN, 2012. Molecular Module: Data Management and Statistical Analysis Plan. Version 1.0. Available at: <http://www.wwarn.org/sites/default/files/MolecularDMSAP.pdf>. Accessed Jan 13, 2014.
57. WWARN, 2012. Statistical Analysis Plan ASAQ/AL Molecular Marker Study Group. Version 1.0. Available at: <http://www.wwarn.org/sites/default/files/ASAQALMolecularMarkerStudyGroupSAP.pdf>. Accessed Jan 13, 2014.
58. Schoenfeld D, 1982. Partial residuals for the proportional hazards regression model. *Biometrika* 69: 239- 241.
59. Choudhury J, 2002. Non-parametric confidence interval estimation for competing risks analysis: application to contraceptive data. *Statistics in medicine* 21: 1129-1144.
60. Ezzet F, van VM, Nosten F, Looareesuwan S, White NJ, 2000. Pharmacokinetics and pharmacodynamics of lumefantrine (benflumetol) in acute falciparum malaria. *Antimicrob.Agents Chemother.* 44: 697-704.
61. Duah N, Mtrevi S, de Souza D, Binnah D, Tamakloe M, Opoku V, Onwona C, Narh C, Quashie N, Abuaku B, Duplessis C, Kronmann K, Koram K, 2013. Increased *pfmdr1* gene copy number and the decline in *pfcr1* and *pfmdr1* resistance alleles in Ghanaian *Plasmodium falciparum* isolates after the change of anti-malarial drug treatment policy. *Malaria Journal* 12: 377.
62. Pickard AL, Wongsrichanalai C, Purfield A, Kamwendo D, Emery K, Zalewski C, Kawamoto F, Miller RS, Meshnick SR, 2003. Resistance to antimalarials in Southeast Asia and genetic polymorphisms in *pfmdr1*. *Antimicrob Agents Chemother* 47: 2418-23.
63. Arie F, Witkowski B, Amaratunga C, Beghain J, Langlois A-C, Khim N, Kim S, Duru V, Bouchier C, Ma L, Lim P, Leang R, Duong S, Sreng S, Suon S, Chuor CM, Bout DM, Menard S, Rogers WO, Genton B, Fandeur T, Miotto O, Ringwald P, Le Bras J, Berry A, Barale J-C, Fairhurst RM, Benoit-Vical F, Mercereau-Puijalon O, Menard D, 2014. A molecular marker of artemisinin-resistant *Plasmodium falciparum* malaria. *Nature* 505: 50-55.
64. Boni MF, Smith DL, Laxminarayan R, 2008. Benefits of using multiple first-line therapies against malaria. *Proc.Natl.Acad.Sci.U.S.A* 105: 14216-14221.
65. Shretta R, 2008. Operational Challenges of Implementing Multiple First-Line Therapies for Malaria in Endemic Countries. Arlington, VA:: Management Sciences for Health.

66. Sutherland CJ, Babiker H, Mackinnon MJ, Ranford-Cartwright L, El Sayed BB, 2011. Rational deployment of antimalarial drugs in Africa: should first-line combination drugs be reserved for paediatric malaria cases? *Parasitology* 138: 1459-68.
67. Eshetu T, Berens-Riha N, Fekadu S, Tadesse Z, Gurkov R, Holscher M, Loscher T, Miranda IB, 2010. Different mutation patterns of *Plasmodium falciparum* among patients in Jimma University Hospital, Ethiopia. *Malar.J.* 9: 226.
68. Eshetu T, Abdo N, Bedru KH, Fekadu S, Wieser A, Pritsch M, Loscher T, Berens-Riha N, 2012. Open-label trial with artemether-lumefantrine against uncomplicated *Plasmodium falciparum* malaria three years after its broad introduction in Jimma Zone, Ethiopia. *Malar J* 11: 240.
69. Borrmann S, Sasi P, Mwai L, Bashraheil M, Abdallah A, Muriithi S, Fruhauf H, Schaub B, Pfeil J, Peshu J, Hanpithakpong W, Rippert A, Juma E, Tsofa B, Mosobo M, Lowe B, Osier F, Fegan G, Lindegardh N, Nzila A, Peshu N, Mackinnon M, Marsh K, 2011. Declining responsiveness of *Plasmodium falciparum* infections to artemisinin-based combination treatments on the Kenyan coast. *PLoS One* 6: e26005.
70. Andriantsoanirina V, Ratsimbaoa A, Bouchier C, Tichit M, Jahevitra M, Rabearimanana S, Raherinjafy R, Mercereau-Puijalon O, Durand R, Menard D, 2010. Chloroquine clinical failures in *P. falciparum* malaria are associated with mutant *Pfmdr-1*, not *Pfcr1* in Madagascar. *PLoS.One.* 5: e13281.
71. Ngasala BE, Malmberg M, Carlsson AM, Ferreira PE, Petzold MG, Blessborn D, Bergqvist Y, Gil JP, Premji Z, Bjorkman A, Mårtensson A, 2011. Efficacy and effectiveness of artemether-lumefantrine after initial and repeated treatment in children <5 years of age with acute uncomplicated *Plasmodium falciparum* malaria in rural Tanzania: a randomized trial. *Clin.Infect.Dis.* 52: 873-882.
72. Kamugisha E, Jing S, Minde M, Kataraihya J, Kongola G, Kironde F, Swedberg G, 2012. Efficacy of artemether-lumefantrine in treatment of malaria among under-fives and prevalence of drug resistance markers in Igombe-Mwanza, north-western Tanzania. *Malar.J.* 11: 58.
73. Sisowath C, Petersen I, Veiga MI, Mårtensson A, Premji Z, Bjorkman A, Fidock DA, Gil JP, 2009. In vivo selection of *Plasmodium falciparum* parasites carrying the chloroquine-susceptible *pfcr1* K76 allele after treatment with artemether-lumefantrine in Africa. *J.Infect.Dis.* 199: 750-757.
74. Baliraine FN, Rosenthal PJ, 2011. Prolonged selection of *pfmdr1* polymorphisms after treatment of falciparum malaria with artemether-lumefantrine in Uganda. *J.Infect.Dis.* 204: 1120-1124.
75. Zongo I, Dorsey G, Rouamba N, Dokomajilar C, Sere Y, Rosenthal PJ, Ouedraogo JB, 2007. Randomized comparison of amodiaquine plus sulfadoxine-pyrimethamine, artemether-lumefantrine, and dihydroartemisinin-piperaquine for the treatment of uncomplicated *Plasmodium falciparum* malaria in Burkina Faso. *Clin.Infect.Dis.* 45: 1453-1461.
76. Ursing J, Kofoed PE, Rodrigues A, Blessborn D, Thoft-Nielsen R, Bjorkman A, Rombo L, 2011. Similar efficacy and tolerability of double-dose chloroquine and artemether-lumefantrine for treatment of *Plasmodium falciparum* infection in Guinea-Bissau: a randomized trial. *J.Infect.Dis.* 203: 109-116.
77. Schramm B, Valeh P, Baudin E, Mazinda CS, Smith R, Pinoges L, Dhorda M, Boum Y, 2nd, Sundaygar T, Zolia YM, Jones JJ, Comte E, Houze P, Jullien V, Carn G, Kiechel JR,

- Ashley EA, Guerin PJ, 2013. Efficacy of artesunate-amodiaquine and artemether-lumefantrine fixed-dose combinations for the treatment of uncomplicated *Plasmodium falciparum* malaria among children aged six to 59 months in Nimba County, Liberia: an open-label randomized non-inferiority trial. *Malar J* 12: 251.
78. Falade CO, Ogundele AO, Yusuf BO, Ademowo OG, Ladipo SM, 2008. High efficacy of two artemisinin-based combinations (artemether-lumefantrine and artesunate plus amodiaquine) for acute uncomplicated malaria in Ibadan, Nigeria. *Trop Med Int Health* 13: 635-43.
79. Wong RP, Karunajeewa H, Mueller I, Siba P, Zimmerman PA, Davis TM, 2011. Molecular assessment of *Plasmodium falciparum* resistance to antimalarial drugs in Papua New Guinea using an extended ligase detection reaction fluorescent microsphere assay. *Antimicrob. Agents Chemother.* 55: 798-805

Tables And Figures

Table 1. Multivariable risk factors for PCR-adjusted recrudescence of artemether-lumefantrine and artesunate-amodiaquine at day 42 based upon previous studies (references will be listed at publication)

Treatment	Variable	Adjusted HR [95% CI]	p-value
AL (N=4,433; 150 recrudescences)			
	Age category: ≥ 12 years (reference)		
	< 1 year	1.96 [0.73 – 5.32]	0.184
	1 to < 5 years	2.05 [1.23 – 3.39]	0.006
	5 to < 12 years	1.23 [0.68 – 2.21]	0.488
	Enrolment parasite density (log-scale)	1.13 [1.05 – 1.23]	0.002
	Lumefantrine dose (mg/kg)	0.99 [0.98 – 1.00]	0.086
ASAQ (N=7,652; 220 recrudescences)			
	Age category: ≥ 12 years (reference)		
	< 1 year	2.20 [1.01 - 4.78]	0.047
	1 to < 5 years	2.27 [1.13 - 4.55]	0.021
	5 to < 12 years	1.51 [0.72 - 3.17]	0.140
	Enrolment parasite density (log-scale)	1.50 [1.16 - 1.93]	0.002
	Amodiaquine dose (mg/kg)	0.92 [0.82 - 1.04]	0.180
	Drug formulation: Fixed dose (reference)		
	Co-blistered	0.98 [0.41 - 2.32]	0.960
	Loose	2.94 [1.58 - 5.48]	0.001

Table 2. PCR-adjusted adequate clinical and parasitological response (ACPR) of artemether-lumefantrine and artesunate-amodiaquine after 42 days of follow up

	AL	ASAQ fixed dose and co-blistered	ASAQ loose
At risk (N)	4,763	1,113	986
ACPR by group (% , [95% CI])			
Age Category			
< 1 year	96.7 [92.7-98.5]	100	85.2 [70.5-93.0]
1-<5 years	93.6 [92.1-94.8]	96.4 [93.2-98.1]	93.8 [90.0-96.2]
5-12 years	96.7 [94.9-97.8]	98.8 [91.6-99.8]	99 [96.1-99.8]
>= 12 years	96.6 [95.2-97.6]	-	-
Region			
Asia/Oceania	97.0 [95.6-97.9]	-	-
East Africa	93.8 [92.4-95.0]	100*	91.2 [88.0-94.7]
West Africa	96.2 [94.6-97.3]	96.9 [94.2-98.3]	99.2 [96.8-99.8]*
Overall	95.3 [94.4-96.0]	97.0 [94.4-98.4]	93.0 [89.2-95.6]

*Followed up to day 28

Table 3. Baseline (pre-treatment) prevalence of genetic markers associated with drug resistance

Markers	Asia/Oceania	East Africa	West Africa
<i>pfprt</i> 76			
Sample size	84	1708	1848
K	3 (4%)	553 (32%)	494 (27%)
K/T	2 (2%)	125 (7%)	249 (13%)
T	79 (94%)	1030 (60%)	1105 (60%)
<i>pfprt</i> 72-76			
Sample size	84	155	84
CVMNK	3 (4%)	37 (24%)	14 (17%)
CVIET	0	117 (75%)	53 (63%)
SVMNT	79 (94%)	0	0
mixed	2 (2%)	1 (1%)	17 (20%)
<i>pfmdr1</i> 86			
Sample size	156	2033	1321
N	97 (62%)	759 (37%)	678 (51%)
N/Y	0	378 (19%)	190 (14%)
Y	59 (38%)	896 (44%)	453 (34%)
<i>pfmdr1</i> 184			
Sample size	158	1275	686
Y	133 (84%)	803 (63%)	287 (42%)
Y/F	7 (4%)	130 (10%)	77 (11%)
F	18 (11%)	342 (27%)	322 (47%)
<i>pfmdr1</i> 1246			
Sample size	64	1017	687
D	54 (84%)	454 (45%)	526 (77%)
D/Y	10 (16%)	309 (30%)	86 (13%)
Y	0	254 (25%)	75 (11%)
<i>pfmdr1</i> 86 + 1246			
Sample size	56	1000	685
N D	2 (3%)	129 (13%)	263 (38%)
N Y	0	9 (1%)	2 (0%)
Y D	47 (84%)	248 (25%)	199 (29%)
Y Y	0	220 (22%)	71 (10%)
mixed	7 (13%)	394 (39%)	150 (22%)
<i>pfmdr1</i> copy number			
Sample size	106	659	0
1	52 (29%)	642 (98%)	0
2	36 (34%)	16 (2%)	0
>2	18 (17%)	1 (0%)	0

Table 4. Multivariable risk factors for PCR-adjusted recrudescence infections of artemether-lumefantrine on day 42

Marker	Variable	Adjusted Hazard Ratio [95% CI]	P-value
<i>pfmdr1</i> 86 (N=2,474; 117 recrudescence infections)*			
	<i>pfmdr1</i> N86 or N/Y:		
	in recrudescence up to day 14	0.90 [0.28 - 2.89]	0.858
	in recrudescence between day 14-28	4.06 [1.94– 8.47]	< 0.001
	in recrudescence after day 28	0.92 [0.46 - 1.81]	0.805
	Enrolment parasite density (log _e -scale)	1.12 [0.97 - 1.30]	0.116
	Age category (reference: < 1 year)		
	1 to < 5 years	1.08 [0.41 - 2.82]	0.882
	5 to < 12 years	0.80 [0.28 – 2.26]	0.671
	>= 12 years	0.74 [0.23 - 2.35]	0.611
	Lumefantrine dose (mg/kg)	0.99 [0.98 - 1.00]	0.219
<i>pfmdr1</i> copy number(N=739; 54 recrudescence infections)			
	<i>pfmdr1</i> copy number > 1:**		
	in recrudescence up to day 14	1.15 [0.11 – 11.95]	0.907
	in recrudescence between day 14-21	2.18 [0.58 – 8.13]	0.247
	in recrudescence after day 21	0.94 [0.29 – 3.11]	0.922
	Region (reference: Africa)		
	Asia/Oceania	11.94 [2.33 – 61.06]	0.003
	Enrolment parasite density (log _e -scale)	1.04 [0.85 - 1.26]	0.707
	Age category (reference: < 5 years)		
	5 to < 12 years	0.86 [0.34 – .21]	0.761
	>= 12 years	0.30 [0.09- 1.07]	0.064
	Lumefantrine dose (mg/kg)	0.98 [0.96 - 1.00]	0.092

*Region not included as a covariate or interaction term with *pfmdr1* 86 genotype because proportional hazards assumption was not met

**Sparse data on *pfmdr1* copy number in Africa prevented the inclusion of region as an interaction term

Table 5. Selection of *pfprt* and *pfmdr1* genotypes after treatment with artemether-lumefantrine and artesunate-amodiaquine

Marker	Genotype	Recurrence		Recrudescence		Re-infection	
		AL	ASAQ	AL	ASAQ	AL	ASAQ
<i>pfprt</i> 76	K --> T ^a	16% (89/571)	10% (25/237)	5% (4/73)	20% (7/35)	17% (82/493)	9% (17/196)
	T --> K	30% (171/571)	8% (18/237)	25% (18/73)	11% (4/35)	31% (152/493)	7% (14/196)
	no change	54% (311/571)	82% (194/237)	70% (51/73)	69% (24/35)	53% (259/493)	84% (165/196)
		<0.001	0.286	0.004 (exact)	0.366	<0.001	0.590
<i>pfmdr1</i> 86	N --> Y	13% (95/692)	27% (92/341)	11% (10/87)	18% (5/28)	14% (85/603)	28% (87/308)
	Y --> N	41% (285/692)	16% (54/341)	36% (31/87)	14% (4/28)	42% (254/603)	16% (49/308)
	no change	45% (312/692)	57% (195/341)	53% (46/87)	68% (19/28)	44% (264/603)	56% (172/308)
		<0.001	0.002	0.001	0.739	<0.001	0.001
<i>pfmdr1</i> 184	Y--> F	25% (73/291)	12% (37/303)	26% (14/55)	12% (3/25)	25% (59/236)	12% (34/273)
	F --> Y	18% (51/291)	17% (50/303)	18% (10/55)	4% (1/25)	17% (41/236)	18% (49/273)
	no change	57% (167/291)	71% (216/303)	56% (31/55)	84% (21/25)	58% (136/236)	70% (190/273)
		0.048	0.163	0.414	0.625	0.072	0.100
<i>pfmdr1</i> 1246	D --> Y	14% (38/273)	32% (102/317)	11% (5/44)	39% (11/28)	15% (33/227)	32% (90/284)
	Y --> D	32% (86/273)	19% (60/317)	30% (13/44)	14% (4/28)	32% (73/227)	20% (56/284)
	no change	54% (149/273)	49% (155/317)	59% (26/44)	46% (13/28)	53% (121/227)	48% (138/284)
		<0.001	0.001	0.059	0.119	<0.001	0.005
<i>pfmdr1</i> copy number	1 --> 2 or more	0.5% (1/247)	--	3% (1/37)	--	0	--
	2 or more --> 1	1% (2/247)	--	0	--	1% (2/210)	--
	no change	98.5% (244/247)	--	97% (36/37)	--	99% (208/210)	--
		1.000 (exact)	--	0.317 (exact)	--	0.500 (exact)	--

Changes in bold indicate statistically significant selection ($p < 0.05$) using McNemar's paired test. Those marked (exact) were tested using the exact distribution for small sample sizes. A small number of recurrent infections (4 for AL and 6 for ASAQ) were not PCR-adjusted and were excluded from the analysis of recrudescence and re-infections.

^a Each category includes all changes from one allele to another. For example, $K \rightarrow T$ includes $K \rightarrow T$, $K \rightarrow K/T$ and $K/T \rightarrow T$ changes





Figure 1. Patient flowchart

A



B








Figure 2. Polymerase chain reaction-adjusted efficacy as assessed by Kaplan-Meier survival estimates for artemether-lumefantrine (AL) by *pfmdr1* genotype of initial parasites. Dotted line indicates WHO-recommended 90% efficacy cutoff for antimalarials. Clinical response of patients with parasites that carry A) *pfmdr1* 86Y (blue) versus 86N or N/Y (red); N = 2,474 patients at risk B) *pfmdr1* copy number > 1 (yellow) versus single copy (green); N = 739 patients at risk.

Figure 3A.










Figure 3A. Cumulative (left panels) and relative (right panels) risks of PCR-adjusted re-infection for baseline *pfprt* and *pfmdr1* genotypes after artemether-lumefantrine treatment, where recrudescent and re-infections with other genotypes were treated as competing events.

Figure 3B.






Figure 3B. Cumulative (left panels) and relative (right panels) risks of PCR-adjusted re-infection for baseline *pfprt* and *pfmdr1* genotypes after artesunate-amodiaquine treatment, where recrudescent and re-infections with other genotypes were treated as competing events